



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C07C 249/12</b>	<b>A1</b>	(11) International Publication Number: <b>WO 98/43948</b>
		(43) International Publication Date: 8 October 1998 (08.10.98)

(21) International Application Number: PCT/HU97/00082

(22) International Filing Date: 5 December 1997 (05.12.97)

(30) Priority Data:  
P 97 00699 3 April 1997 (03.04.97) HU(71) Applicant (for all designated States except US): BIOREX  
KUTATÓ ÉS FEJLESZTŐ RT. [HU/HU]; Szabadságpuszta,  
H-82001 Veszprém (HU).




(72) Inventors; and

(75) Inventors/Applicants (for US only): ÜRÖGDI, László  
[HU/HU]; Teleki u. 80, H-1184 Budapest (HU).  
BARABÁS, Mihály [HU/HU]; Egry J. u. 36, H-1111  
Budapest (HU). MÁRVÁNYOS, Ede [HU/HU]; Béke tér  
5, H-1139 Budapest (HU). TÖRÖK, Magdolna [HU/HU];  
Zöldfa u. 172, H-4700 Mátészalka (HU). CSÁKAI, Zita  
[HU/HU]; Petőfi ltp. B/19, H-6090 Kunszentmiklós (HU).(74) Agent: S.B.G. AND K. PATENT AND LAW OFFICE;  
Andrássy út 113, H-1062 Budapest (HU).(81) Designated States: AL, AU, BA, BB, BG, BR, CA, CN, CU,  
CZ, EE, GE, HU, IS, JP, KP, KR, LC, LK, LR, LT, LV,  
MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL,  
TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, KE,  
LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY,  
KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH,  
DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT,  
SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML,  
MR, NE, SN, TD, TG).

Published

With international search report.

(54) Title: PROCESS FOR PREPARING O-(3-AMINO-2-HYDROXY-PROPYL)-HYDROXYMIC ACID HALIDES



## (57) Abstract

The present invention relates to a novel process for preparing O-(3-amino-2-hydroxy-propyl)-hydroxymic acid halides of formula (I) by reacting a carboxamide oxime of formula (II) wherein R<sup>1</sup> is as specified above with reactive 3-amino-2-hydroxy-propane derivative, diazotizing the O-substituted carboxamide oxime thus obtained with sodium nitrite in the presence of hydrogen halide, decomposing the diazonium salt and if desired, separating the optically active enantiomers and/or reacting the resulting base with an organic or mineral acid, wherein the carboxamide oxime of formula (II) is reacted with a 3-hydroxy azetidinium salt of formula (III) wherein R<sup>2</sup> and R<sup>3</sup> are as defined above and Y<sup>-</sup> is a salt forming anion, in a lower alcoholic, preferably ethanolic medium optionally containing water and made alkaline with an alkali hydroxide, and before diazotizing the O-substituted carboxamide oxime intermediate obtained, the reaction mixture is neutralised and the organic solvent is removed. The process according to the invention provides the compounds of formula (I) with a higher yield compared to the prior art processes.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

5 PROCESS FOR PREPARING  
O-(3-AMINO-2-HYDROXY-PROPYL)-HYDROXYMIC ACID HALIDES

### Technical field

The present invention relates to a novel process for preparing O-(3-amino-2-hydroxy-10 propyl)-hydroxymic acid halides of the formula (I) wherein

R<sup>1</sup> is phenyl, pyridyl or thienyl or substituted phenyl, wherein the one or more substituent(s) may be halo and/or haloalkyl and/or nitro,

X is halo,

15  $R^2$  and  $R^3$  are independently from each other straight or branched lower alkyl or  $R^2$  and  $R^3$  together with the nitrogen connecting thereto form a saturated 5 to 7-membered heterocyclic group which may contain additional hetero atom and may be substituted.

The invention also relates to a process for preparing the acid addition salts and optically active forms of the above compounds.

### Background art

O-(3-amino-2-hydroxy-propyl)-hydroxymic acid halides of the general formula (I) are well known as active substances in the treatment of pathological changes in the vascular system connected with diabetes mellitus, especially with diabetic angiopathy.

25 These compounds are particularly described e.g. in the Hungarian patent No. 207.988.

O-(3-amino-2-hydroxy-propyl)-hydroxymic acid halides of the general formula (I) can be prepared in many different ways some of them being also described in the Hungarian patent No. 207.988.

Although the known synthesis routes for manufacture are suitable allow the preparation of the compounds of formula (I), they are not sufficient for the preparation of the said compounds in industrial scale. The disadvantage thereof is that they need reagents which are difficult to handle or prepare or comprise unfavourable reactions with non-satisfactory yields due to the possibility of side reactions. The urging need for the compounds of formula (I) requires a novel process which is secure, has a satisfactory yield and can be carried out under industrial conditions.

#### Disclosure of invention

10 The present invention aims to provide a process for preparing O-(3-amino-2-hydroxy-propyl)-hydroxymic acid halides in industrial scale.

The present invention provides an industrially applicable process for preparing the compounds of the formula (I) by

- i) reacting an amidoxime compound of the formula II with a 3-hydroxy-acetidinium salt wherein
- 15  $R^2$  and  $R^3$  have the meaning as specified above and  
 $Y^-$  is a salt forming anion  
in a basic-alcoholic medium,
- ii) neutralizing the mixture and removing the organic solvent,
- 20 iii) reacting the residue with sodium nitrite in aqueous medium in the presence of hydrochloric acid
- iv) decomposing the diazonium salt thus obtained and
- v) isolating the crude product of the formula (I) from the mixture.

Reactions of amidoxime compounds of the formula II and suitably substituted 3-amino-2-hydroxy-propane derivatives (usually 1-halo- or 1,2-epoxy derivatives) are described e.g. in the Hungarian patent No. 177.578. However, the 3-hydroxy azetidinium salts of the formula (III) are more suitable reagents than the 1-halo- or 1,2-epoxy derivatives used in the known reactions. Namely, compounds of the formula (III) are solid materials which can easily be prepared, isolated and stored unlike the reagents

30 formerly used which are difficult to isolate and handle and are generally liquid

materials. The use thereof became known from Hungarian patent specification No. 207.305, but in the process described therein neither these nor the other two reagents are directly reacted with the compounds of the formula (II), but with an amidoxime complex prepared therefrom with alkali hydroxide or alkali and dimethyl formamide alcoholate or 1,3-dimethyl-2-imidazolidinone in a medium containing dimethyl formamide. Thus, O-substituted amidoxime derivatives were isolated which, however appear only as non-isolated intermediates in the reaction sequence according to the present invention.

Based on our observations efforts were made to eliminate the tecnologically difficult complex forming from the process and to avoid the use of dimethyl formamide as solvent. Dimethyl formamide is hazardous for health as it causes cancer, furthermore it is difficult to regenerate and purify and extremely difficult to make water-free. This is especially important as dimethyl formamide impurity must be minimized as pointed out in Hungarian patent No. 207.305. Moreover, it is preferable that the solvent contains only very small quantity of water, practically less than 1% to achieve a proper yield. An additional disadvantage of the use of dimethyl formamide is that it decomposes when exposed to light thus becoming contaminated by the toxic compounds envolved. It has been found that the reaction can safely and easily be carried out by reacting the compounds of the formulae (II) and (III) directly in a basic alcoholic medium which may also contain water. In respect of the outcome of the synthesis it is very useful that the O-substituted carboxamide oxime intermediate is not isolated from the reaction mixture, but directly reacted further after neutralizing the mixture and removing the organic solvent. It has also been found that the side products formed during the contracted steps can all be removed by one suitable isolation step and thus, the synthesis is appropriate for manufacturing the product in the desired purity.

Based on these observations, the invention provides a process for preparing compounds of the formula (I), wherein

$R^1$  is phenyl, pyridyl or thienyl or substituted phenyl, wherein the one or more substituents may be halo and/or haloalkyl and/or nitro,

X is halo,

$R^2$  and  $R^3$  are independently from each other straight or branched lower alkyl or  
 $R^2$  and  $R^3$  together with the nitrogen connecting thereto form a 5 to 7-membered  
saturated heterocyclic group which may contain additional hetero atom and may  
be substituted.

and the acid addition salts and optically active forms thereof by reacting a carboxamide  
oxime of the formula (II) wherein the meaning of  $R^1$  is as specified above with a  
reactive 3-amino-2-hydroxy-propane derivate, diazotizing the resulting O-substituted  
carboxamide oxime with sodium nitrite in the presence of hydrohalide, decomposing  
the diazonium salt, isolating the product obtained and, if desired, separating the  
optically active enantiomers and/or reacting the resulting base with an organic or  
mineral acid; which comprises reacting the carboxamide oxime of the formula (II) with  
a 3-hydroxy acetidinium salt of the formula (III) wherein  $R^2$  and  $R^3$  have the meaning  
as specified above and  $Y^-$  is a salt forming anion in a lower alcoholic preferably  
ethanolic medium made alkaline with an alkali hydroxide, the said medium optionally  
containing water while neutralizing the reaction mixture and removing the organic  
solvent therefrom before diazotizing the resulting O-substituted carboxamide oxime  
intermediate.

#### Best mode for carrying out the invention

A preferred mode of carrying out the process according to the invention is as follows:

The carboxamid oxime of the formula (II) and the 3-hydroxy azetidinium salt of  
the formula (III) are reacted at stoichiometric ratio, however, it may be advantageous  
to apply the compound of the formula (III) in a slight excess. The reaction can be  
carried out with any order of addition of the reagents, preferably the compound of the  
formula (II) is added to the basic-alcoholic solution of the compound of the formula  
(III). As solvent preferably a  $C_{1-4}$  alkanol, preferably ethanol is used and the reaction is  
carried out preferably with heating, most preferably at the boiling point of the solvent.  
After termination of the reaction, the mixture is cooled and neutralized with a mineral  
acid, preferably hydrochloric acid, and the alcohol is distilled off suitably at lower

pressure. After removing the solvent the reaction mixture is diluted with water, the concentrated hydrochloric acid necessary for diazotizing is added, cooled to the diazotizing temperature and is diazotized by the addition of the sodium nitrite under cooling at a temperature of 0 to + 5 C°. The diazonium salt decomposes in situ into the  
5 corresponding hydroximoyl-halide derivative. To isolate the crude product, the reaction mixture is made alkaline with an inorganic alkali compound, extracted with an organic solvent non-miscible with water, preferably ethyl acetate, the extract is dried and concentrated, or directly an acid addition salt is formed from the product by adding a suitable acid to the mixture and separating the acid addition salt by filtration. The  
10 crude product can be purified by recrystallization or by any other way known in the art.

By using the above process, a properly pure product can be prepared with an economically satisfactory yield.

The advantage of the present invention lies in the fact that it makes possible to produce O-(3-amino-2-hydroxy-propyl)-hydroxime acid halides by a safe and simple  
15 method also under industrial conditions.

The invention is further illustrated in the following examples:

#### Example 1.

N-[2-hydroxy-3-(1-piperidiny)-propoxy]-3-pyridine-carboximidoyl chloride (Z)-2-  
20 butenedioate

50.4 kg 2-hydroxy-4-azoniaspiro-[3,5]-nonane chloride was dissolved in and 28 l of water under stirring. To the solution 11,4 kg sodium hydroxide was added and the resulting milk-like mixture was stirred for an additional hour. While stirring, 420 l of  
25 ethanol and 35 kg 3-pyridine-carboxamide oxime were added thereto and the mixture was heated under reflux for 1.5 hours followed by cooling the reaction mixture. 270-290 l of alcohol were distilled off and 110 l of deionized water and 45.5 l of concentrated hydrochloric acid were added followed by distilling off the remaining ethanol. To the oily residue 160 l of concentrated hydrochloric acid was added under  
30 cooling so that the temperature remained under 30 C°. The solution was then cooled to

0 C° and for the diazotization a mixture of 17.7 kg sodium nitrite and 60 l of deionized water was added under permanent stirring and cooling while maintaining the temperature of the reaction mixture between 0 and +5 C°. After the addition, the mixture was stirred for an additional hour at this temperature and for the  
5 decomposition of the nitrite excess 1.5 kg urea was added. After full decomposition of the nitrite (appr. 1.5 hours) to the reaction mixture 350 l ethyl acetate was added followed by alkalifying by the addition of 150-200 l concentrated sodium hydroxide under stirring and intensive cooling. The layers were separated, the organic phase washed with 2 x 70 l of water and dried over 15 kg anhydrous Na<sub>2</sub>SO<sub>4</sub>. The drying  
10 agent was filtered off, washed with 20 l ethyl acetate, the organic layers combined and the quantity of the N-[2-hydroxy-3-(piperidine-1-yl)-propoxy-3-pyridine-carboximidoyl chloride base was determined. Maleic acid in calculated amount (21-22 kg) was added and the mixture was stirred for 4 hours. The product was separated in centrifuge, washed with 30 l acetone and the resulting crude product was dissolved in  
15 70 l warm acetone and recrystallized. The product was separated in centrifuge and washed with 30 l acetone. After recrystallization, 50-55 kg N-[2-hydroxy-3-(1-piperidinyl)-propoxy]-3-pyridine-carboximidoyl chloride (Z)-2-butenedioate (1:1) was obtained as pink beige crystals. (m.p. 123-124 C°, acetone, yield: 53%).

20 IR (v, KBr/cm<sup>-1</sup>): 3350, 2941, 1580, 1480, 1350, 1022, 982, 867, 702.

<sup>1</sup>H-NMR (250 MHz, DMSO-d<sub>6</sub>; ref.: DMSO-d<sub>6</sub>=2,5.8(ppm): 9.00 (1H, s); 8.74 (1H, d); 8.18 (1H, d), 7.56(1H, dd); 6.03 (2H, s); 5.85-6.00 (1H, s/br); 4.21-4.37 (3H, m); 3.2-3.33 (2H, m); 2.49-2.55 (4H, m); 1.54-1.77 (6H, m).

25

<sup>13</sup>C-NMR (63 MHz, solvent: DMSO-d<sub>6</sub>; ref.: DMSO-d<sub>6</sub>=39.3.8(ppm): 167.0 (COOH); 151.4, 127.9, 134.3, 123.5, 147.2 (pyridine 2-3-4-5-6); 135.4 (CH=CH); 134.9 (C/Cl=NO); 77.2 (NOCH<sub>2</sub>); 63.5 (CHOH, 58.3 (NCH<sub>2</sub>); 52.9, 22.1, 21.2 (piperidine).



According to Example 1 the following compounds were prepared:

Example 2.

N-[2-hydroxy-3-(1-piperidinyl)-propoxy]-benzimidoyl chloride hydrochloride

5 (m.p.: 140-144 C°, isopropanol, yield: 66%)

IR (KBr): 3234, 2951, 1504, 1448, 1389, 1289, 1119, 1059, 972, 768, 690.

Example 3.

N-{2-hydroxy-3-[1-(4-methyl)-piperazinyl]-propoxy}-3-pyridine-carboximidoyl

10 chloride (Z)-2-butenedioate (1:2)

(m.p.: 174-175 C°, ethanol, yield: 48%)

IR (KBr): 3207, 1693, 1578, 1456, 1358, 1304, 1020, 974, 864, 702.

Example 4.

15 N-[2-hydroxy-3-(diethylamino)-propoxy]-3-pyridine-carboximidoyl chloride hydrochloride

(m.p.: 118-119 C°, acetone, yield: 67%)

IR (KBr): 3425, 3289, 2951, 2667, 1818, 1443, 1337, 1238, 1178, 1115, 1078, 1049, 997, 910, 804, 781, 696, 683 cm<sup>-1</sup>

20

Example 5.

N-[2-hydroxy-3-(4-morpholinyl)-propoxy]-3-pyridine-carboximidoyl chloride (Z)-2-butenedioate

(m.p.: 137-138 C°, isopropanol, yield: 52%)

25 IR (KBr): 3310, 1580, 1483, 1464, 1443, 1354, 1072, 1024, 982

## Example 6.

N-[2-hydroxy-3-(1-piperidiny)-propoxy]-2-tiophene-carboximidoyl chloride hydrochloride

(m.p.: 115-123 C°, isopropanol-hexane, yield: 38%)

5

<sup>1</sup>H-NMR (250 MHz, DMSO-d<sub>6</sub>; ref.: DMSO-d<sub>6</sub>=2,5δ(ppm): 10.2 (1H, s/br); 7.81, 7.63, 7.20 (1H, 1H, 1H, d, d, dd); 5.98 (1H, s/b); 4.42 (1H, s/b), 4.35 (2H, d); 3.60-2.90 (6H, m); 1.95-1.60 (4H, m); 1.45-1.20 (2H, m).

10 <sup>13</sup>C-NMR (63 MHz, DMSO-d<sub>6</sub>; ref.: DMSO-d<sub>6</sub>=39.3δ(ppm): 133.8 [C(Cl)=N]; 132.1, 130.3, 130.1, 127.6 (tiophene 3-2-5-4); 76.8 (NOCH<sub>2</sub>); 63.2 (CHOH); 58.5 (CH<sub>2</sub>N); 53.3, 51.8 (piperidine 2 x NCH<sub>2</sub>) 22.0, 21.9, 21.0 (piperidine 3 x CH<sub>2</sub>).

## Example 7

15 N-[2-hydroxy-3-(1-piperidiny)-propoxy]-2-trifluoromethyl benzimidoyl chloride hydrochloride

(m.p.: 119-123 C°, ethylacetate, yield: 30%)

IR (KBr): 3366, 2937, 2854, 2737, 2673, 2538, 1616, 1570, 1439, 1404, 1337, 1290, 1236, 1199, 1165, 1129, 1101, 1074, 1030, 984, 972, 933, 901, 829, 804, 788, 717,  
20 699, 685, 646 cm<sup>-1</sup>.

## Example 8

N-[2-hydroxy-3-(1-piperidiny)-propoxy]-2'-nitrobenzimidoyl chloride hydrochloride

(m.p.: 159-162 C°, isopropanol, yield: 43%)

25 IR (KBr): 3298, 2983, 2932, 2746, 1593, 1574, 1535, 1445, 1391, 1354, 1317, 1288, 1242, 1198, 1117, 1092, 1069, 1020, 968, 947, 914, 852, 793, 756, 708, 577 cm<sup>-1</sup>.

## Example 9

(+) N-[2-hydroxy-3-(1-piperidin-1-yl)-propoxy]-3-piridine-carboximidoyl chloride (Z)-2-butenedioate (1:1)

5 2-hydroxy-4-azoniaspiro[3.5]nonane chloride and 3-piridine-carboxamide oxime were reacted according to Example 1 following the reaction steps up to separating the N-[2-hydroxy-3-(1-piperidin-1-yl)-propoxy]-3-piridine-carboximidoyl chloride with ethyl acetate. 15 g (50 mmole) N-[2-hydroxy-3-(1-piperidin-1-yl)-propoxy]-3-piridine-carboximidoyl-chloride in ethyl acetate was added dropwise to a mixed anhydride  
10 prepared from 13.52 g (50 mmole) N-(t-butoxycarbonyl)-L-phenylalanine and 5,0 ml ethyl chloroformate in dichlorometane by a method known per se and the mixture was stirred for an hour at room temperature. To isolate the ester thus obtained the solution was extracted with 2 x 200 ml aqueous acetic acid solution (10%) and 1 x 200 ml water, the organic layer was dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and concentrated. The oily  
15 residue was dissolved in 140 ml acetone and to the solution 3,0 g maleic acid was added. Thus, 5.2 g (7.8 mmole, 16%) (-)-N-[2-(N'-BOC-/L/-phenylalanyloxy)-3-(1-piperidinyl)-propoxy]-3-piridine-carboximidoyl chloride (Z)-2-butenedioate (1:1) salt (m.p.: 146.5-148 C°) was obtained.

5.2 g of the salt prepared as above was boiled in methanol for 1 hour. The solution was  
20 distilled to dryness and the residue was crystallized from 50 ml ethyl acetate giving 3.18 g (98%) (+) N-[2-hydroxy-3-(1-piperidin-1-yl)-propoxy]-3-piridine-carboximidoyl chloride (Z)-2-butenedioate (1:1) salt (m.p.: 136-137 C°). The IR and NMR spectrum of the compound corresponded to those of the racemic compound. According to chiral shift spectroscopy the compound was a homogenous enantiomer.  
25 The (-) isomer could be prepared in an analogous way, but using N-(t-butoxycarbonyl)-/D/-phenylalanine as reagent.

The method according to the invention was compared with the method described in the prior art mentioned above. 3-piridine-carboxamide oxime was reacted with 3-  
30 piperidino-2-hydroxy-1-chloropropane prepared according to the method described in

Hungarian patent No. 177.578 in absolute alcoholic medium. After termination of the reaction the solution was made alkaline, the product was extracted with benzene and from the base dihydrochloride was formed with gaseous hydrochloric acid. The O-(3-piperidino-2-hydroxy-1-propyl)-3-pyridine-carboxamide oxime hydrochloride thus  
5 obtained was diazotized according to the method described in Hungarian Patent No. 207.988, the diazonium salt was decomposed and the resulting product was reacted with maleic acid giving the product according to example 1. The final yield of the process based on the starting product was 38 % while the same in Example 1 was 53%, which raised up to 60 % during commercial production.

10 It can be established that the process according to the invention provides the compounds of the formula (I) with a higher yield compared to the prior art processes. A further advantage of the process according to the invention is the possibility to spare solvent. For preparing 1 kg product according to the process described in the present invention only 17 kg solvent was needed while the same according to the formerly  
15 known processes amounted to 40 kg. A further advantage of the process according to the invention in the industrial scale is that the technology time needed for the preparation of the compounds of the formula (I) is shorter. To produce 1 batch of the product related to 3 m<sup>3</sup> reactor volume according to the invention needed 4 consecutive shifts while the prior art processes needed 8 shifts.

20 Summarized, the process according to the present invention provides a method to prepare the O-(3-amino-2-hydroxy-propyl)-hydroxamic acid halides with a higher yield and with substantially reduced technological costs than the processes previously known.

## CLAIM

Process for preparing O-(3-amino-2-hydroxy-propyl)-hydroxamic acid halides of the formula I wherein

5         $R^1$  is phenyl, or pyridyl or thienyl or substituted phenyl, wherein the one or more substituent(s) may be halo and/or haloalkyl and/or nitro,


          X is halo,

$R^2$  and  $R^3$  are independently from each other straight or branched lower alkyl or

10         $R^2$  and  $R^3$  together with the nitrogen connecting thereto form a 5 to 7-membered saturated heterocyclic group which may contain additional hetero atom and may be substituted,

and the acid addition salts and optically active forms thereof by reacting a carboxamide oxime of the formula (II) wherein  $R^1$  is as specified above with reactive 3-amino-2-hydroxy-propane derivative, diazotizing the O-substituted carboxamide oxime thus  
15        obtained with sodium nitrite in the presence of hydrogen halide, decomposing the diazonium salt and if desired, separating the optically active enantiomers and/or reacting the resulting base with an organic or mineral acid, wherein the carboxamide oxime of the formula (II) is reacted with a 3-hydroxy azetidinium salt of the formula (III) wherein  $R^2$  and  $R^3$  are as defined above and  $Y^-$  is a salt forming anion, in a lower  
20        alcoholic, preferably ethanolic medium optionally containing water and made alkaline with an alkali hydroxide, and before diazotizing the O-substituted carboxamide oxime intermediate obtained, the reaction mixture is neutralized and the organic solvent is removed.

1/1



## INTERNATIONAL SEARCH REPORT

International Application No  
PCT/HU 97/00082A. CLASSIFICATION OF SUBJECT MATTER  
IPC 6 C07C249/12

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 6 C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 90 04584 A (BIOREX KUTATO FEJLESZTOE KFT) 3 May 1990 see page 4, line 22 - line 24; examples 3-9	1
Y	& HU 207 988 B (...) cited in the application	1
Y	WO 90 08131 A (CHINOIN GYOGYSZER ES VEGYESZET) 26 July 1990 see page 2, line 23 - line 30; claims 1,7; example 10	1
	--- -/--	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*&\* document member of the same patent family

Date of the actual completion of the international search

10 February 1998

Date of mailing of the international search report

06.03.98

Name and mailing address of the ISA  
European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Janus, S

# INTERNATIONAL SEARCH REPORT

Int. Application No  
PCT/HU 97/00082

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>DATABASE WPI Week 9322 Derwent Publications Ltd., London, GB; AN 93-177741 XP002055093 &amp; HU 207 305 B (CHINOIN GYOGYSZER ES VEGYESZETI) , 29 March 1993 cited in the application see compounds of formula (V)</p>	1
A	<p>GEARTNER, R.V.: "Ring-opening alkylations of 1,1-dialkyl-3-substituted azetidinium cations. Substituent entropy-controlled strained ring-chain equilibria" JOURNAL OF ORGANIC CHEMISTRY, vol. 33, no. 1, 1968, pages 523-530, XP002055092 see page 528, column 2, line 54 - line 77 see page 529, column 1, line 56 - line 66 see page 529, column 2, line 12 - line 31</p>	1



# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/HU 97/00082

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9004584 A	03-05-90	AU 620460 B	20-02-92
		AU 4418689 A	14-05-90
		CA 2000830 A,C	20-04-90
		DE 68913737 D	14-04-94
		DE 68913737 T	28-07-94
		DK 149790 A	19-06-90
		EP 0417210 A	20-03-91
		FI 93214 B	30-11-94
		GR 89100669 A	29-11-90
		HU 9500117 A	28-06-95
		IE 65113 B	04-10-95
		IL 92000 A	24-06-94
		JP 8019078 B	28-02-96
		JP 3502931 T	04-07-91
		NO 178148 B	23-10-95
		PL 164547 B	31-08-94
		PT 92041 B	18-07-95
		RU 2093508 C	20-10-97
		US 5328906 A	12-07-94
		US 5147879 A	15-09-92
		US 5296606 A	22-03-94
WO 9008131 A	26-07-90	AT 132485 T	15-01-96
		AU 639223 B	22-07-93
		AU 4823790 A	13-08-90
		BG 60267 B	31-03-94
		CA 2024808 A	11-07-90
		DE 69024546 D	15-02-96
		DE 69024546 T	29-08-96
		EP 0406386 A	09-01-91
		ES 2082847 T	01-04-96
		HU 9500220 A	28-09-95
		JP 3504016 T	05-09-91
		NO 177821 B	21-08-95
		RU 2074854 C	10-03-97
		US 5278309 A	11-01-94
		US 5239077 A	24-08-93